

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/56983 A2

- (51) International Patent Classification⁷: C07D
- (21) International Application Number: PCT/US01/03336
- (22) International Filing Date: 1 February 2001 (01.02.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/179,505 1 February 2000 (01.02.2000) US
- (71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, OH
45202 (US).
- (72) Inventors: **CAZER, Fredrick, Dana**; 220 Reese Rd.,
Earlville, NY 13312 (US). **PERRY, Gregory, Eugene**;
8 Gerald St., Norwich, NY 13815 (US). **BILLINGS,**
Dennis, Michael; 142 Warner Road, Norwich, NY 13815
(US). **REDMAN-FUREY, Nancy, Lee**; 786 Beaver
Meadow Rd., Smyrna, NY 13464 (US).
- (74) Agents: **REED, T., David et al.**; The Procter & Gam-
ble Company, 5299 Spring Grove Avenue, Cincinnati, OH
45217-1087 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility
model), DK, DK (utility model), DM, DZ, EE, EE (utility
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK
(utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SELECTIVE CRYSTALLIZATION OF 3-PYRIDYL-1-HYDROXYETHYLIDENE-1,1-BISPHOSPHONIC ACID
SODIUM AS THE HEMIPENTAHYDRATE OR MONOHYDRATE

(57) Abstract: The present invention discloses 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium hemipentahydrate and monohydrate, methods of preparing the hemipentahydrate or monohydrate through control of the nucleation temperature and rate of crystallization and pharmaceutical compositions containing one or both of the hydrate forms

WO 01/56983 A2

SELECTIVE CRYSTALLIZATION OF 3-PYRIDYL-1-HYDROXYETHYLIDENE-1,1-BISPHOSPHONIC ACID SODIUM AS THE HEMIPENTAHYDRATE OR MONOHYDRATE

5

10

TECHNICAL FIELD

The present invention relates to 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium hemipentahydrate and monohydrate, compositions containing said hemipentahydrate and/or monohydrate and methods of selective crystallization of the hemipentahydrate or monohydrate.

15

CROSS REFERENCE

This application claims priority under Title 35, United States Code 119(e) from Provisional Application Serial No. 60/179,505, filed February 1, 2000.

BACKGROUND OF THE INVENTION

Bisphosphonates such as 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid (RISEDRONATE) have been proposed for use in the treatment of diseases of bone and calcium metabolism. Such diseases include osteoporosis, hyperparathyroidism, hypercalcemia of malignancy, osteolytic bone metastases, myositis ossificans progressiva, calcinosis universalis, arthritis, neuritis, bursitis, tendonitis and other inflammatory conditions. Paget's disease and heterotrophic ossification are currently successfully treated with both EHDP (ethane-1-hydroxy-1,1-diphosphonic acid) and RISEDRONATE.

25

The bisphosphonates tend to inhibit the resorption of bone tissue, which is beneficial to patients suffering from excessive bone loss. However, in spite of certain analogies in biological activity, all bisphosphonates do not exhibit the same degree of biological activity. Some bisphosphonates have serious drawbacks with respect to the degree of toxicity in animals and the tolerability or negative side effects in humans. The salt and hydrate forms of bisphosphonates alter both their solubility and their bioavailability.

30

It is known in the literature that some bisphosphonic acids and their salts are capable of forming hydrates, risedronate sodium exists in three hydration states: mono, hemipenta and anhydrous. Crystallization procedures which selectively yield the hemipentahydrate form or the

monohydrate form are desirable. This application describes the hemipentahydrate and the monohydrate crystal forms, the compositions containing said hemipentahydrate and monohydrate crystal forms and selective formation of these crystals.

5

SUMMARY OF THE INVENTION


The present invention discloses 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium hemipentahydrate and monohydrate, compositions containing said hemipentahydrate and/or monohydrate and methods of selective crystallization of the hemipentahydrate or monohydrate. The temperature of nucleation and the rate of crystallization are the critical variables that control the ratio of hydrates formed.

10

DETAILED DESCRIPTION OF THE PRESENT INVENTION

The present invention is directed to a 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium hemipentahydrate and monohydrate and compositions containing said hemipentahydrate and monohydrate. A novel process is also disclosed for the selective crystallization of the geminal bisphosphonate, risedronate sodium, 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium as the hemipentahydrate and as the monohydrate.

15



20

Risedronate Sodium

Risedronate sodium, the mono sodium salt of risedronate, exists in three crystalline hydration states: anhydrous, mono and hemipentahydrate. The monohydrate and hemipentahydrate are preferred.

25

The hemipentahydrate is the thermodynamically preferred crystalline form under typical processing conditions based on the observation that monohydrate crystals converted to the hemipentahydrate form.

The monohydrate is by weight of water from about 5.0% to about 7.1% , more preferably from about 5.6% to about 6.5% and most preferably about 5.6%. The monohydrate is further characterized by single crystal X-ray crystallography, and thermogravimetric analysis. The monohydrate form also exhibits identifiable signatures when examined by X-ray powder
5 diffraction, differential scanning calorimetry, Fourier transform infrared spectroscopy or near infrared spectroscopy.

The hemipentahydrate is by weight of water from about 11.9% to about 13.9%, more preferably from about 12.5% to about 13.2% and most preferably about 12.9%. The hemipentahydrate is further characterized by single crystal X-ray crystallography, and
10 thermogravimetric analysis. The hemipentahydrate form also exhibits identifiable signatures when examined by X-ray powder diffraction, differential scanning calorimetry, Fourier transform infrared spectroscopy or near infrared spectroscopy.

The temperature of nucleation and the rate of crystallization are the critical variables that control the ratio of hydrates formed. The nucleation temperature can be controlled by controlling
15 the ratio of water to solute, the solution temperature, and the ratio of organic solvent to water.

The risedronate sodium hemipentahydrate is the thermodynamically favored form under the typical processing conditions described. Compositions comprise by weight of risedronate sodium hydrate from about 50% to about 100%, more preferably from about 80% to about 100%, most preferably from about 90% to about 100% risedronate sodium hemipentahydrate and from
20 about 50% to about 0%, more preferably from about 20% to about 0%, most preferably from about 10% to about 0% risedronate sodium monohydrate.

Altering the processing conditions as described can selectively produce the monohydrate crystal form. Compositions comprise by weight of risedronate sodium hydrate from about 50% to about 99%, more preferably from about 80% to about 99%, most preferably from about 95% to
25 about 99% risedronate sodium monohydrate and from about 50% to about 1%, more preferably from about 20% to about 1%, most preferably from about 5% to about 1% risedronate sodium hemipentahydrate.


The invention further comprises pharmaceutical compositions containing the hemipentahydrate and monohydrate compounds.

30

Definitions and Usage of Terms

The following is a list of definitions for terms used herein:

The term "risedronate", as used herein, denotes 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid and has the following structure:



The compound risedronate is further described in U.S. Patent 5,583,122, Benedict et al., assigned to the Procter & Gamble Co., issued December 10, 1996, and "An American
 5 Conference, Bisphosphonates: Current Status and future Prospects," The Royal College of Physicians, London, England, May 21-22, 1990, organized by IBC Technical Services, both references hereby are incorporated by reference.

As used herein, "solvent", is a substance capable of dissolving another substance to form a uniform solution. The solvent may either be polar or non-polar. Solvents are selected from the
 10 group consisting of alcohols, esters, ethers, ketones, amides, and nitriles. Most preferred is isopropanol.

The Process

The process according to the present invention is characterized in that the process described herein is readily adapted to industrial production. The following non-limiting examples
 15 illustrate the processes of the present invention.

The extent of hydration 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium can be controlled by varying the crystallization parameters to control the temperature of nucleation and rate of crystallization. The ratio of hemipentahydrate to monohydrate crystal forms in the product can be effectively controlled by varying the water to 3-pyridyl-1-
 20 hydroxyethylidene-1,1-bisphosphonic acid sodium ratio and the isopropanol to water ratio as well as the temperature (see below).

General Procedure

An aqueous solution of 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium at
 25 0-75 °C, preferably 25-75 °C, more preferably 45-75 °C

will selectively yield either the monohydrate or the hemipentahydrate crystal forms depending upon the conditions of crystallization. The temperature of nucleation and the rate of crystallization governs the hydrate, varying the ratio of water: isopropanol and varying temperature and cooling ramp of the aqueous solution control the ratios of the hydration states formed.

Table 1 shows eight examples of reaction conditions which selectively produce 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium containing varying hemipenta to monohydrate ratios. The theoretical moisture level for the monohydrate is 5.6% and for the hemipentahydrate is 12.9%.

5

Table 1

Example	Weight of Water (X = weight of compound 2)	Weight of isopropanol	IPA Addition Temp (°C)	Cooling Ramp (° C)	% H2O KF
1	8.72X	2.94X	70	70 hold	5.7
2	22X	160X@0 °C	70	70-0 in 2 min. Quench	5.9
3	7.2X	1.08X	75	75 - 60 in 4 hrs - 25 in 2 hrs.	9.6
4	9X	1.26X	75	75 - 60 in 4 hrs 60 - 25 in 2 hrs	11.4
5	8.2X	0.9X	60	60 hold 4 hrs. 60 - 25 in 2 hrs.	12.3
6	9.5X	1.05X	60	60 - 25 in 2 hrs	13.0
7	8.2X	1.39X	60	60 hold 4 hrs. 60 - 25 in 2 hrs.	13.0
8	8.2X	1.15X	60	60 - 25 in 2 hrs	13.1

Example 1. Hemipentahydrate

Conditions leading to nucleation between 25-70 °C, preferably 50-70 °C with a cooling
 10 ramp of 0.1-5 °C per minute, preferably 0.1-2 °C per minute produce the hemipentahydrate. The
 hemipentahydrate is formed by suspending 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic
 acid in water at about 60 °C, adjusting the pH to 4.7-5.0 with sodium hydroxide, adding
 isopropanol to the resulting solution, cooling the suspension and collecting the product by
 filtration.

15

Example 2. Monohydrate

Conditions leading to nucleation above 45°C, preferably above 55°C with appropriate time at temperature and without a cooling ramp or with a rapid quench produce the monohydrate. Maintaining the aqueous solution of 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium at 70°C, the monohydrate slowly crystallizes and is isolated by filtration of the crystals from the hot solution. Conversely, quenching of the 70°C aqueous solution directly into 0°C isopropanol also yields the crystalline monohydrate.

Compositions

The compounds made herein may be used in pharmaceutical compositions. The term "pharmaceutical composition" means a dosage form comprised of a safe and effective amount of an active ingredient and pharmaceutically-acceptable excipients. The pharmaceutical compositions described herein are comprised of from about, 0.1% to about 99%, preferably from about 0.5% to about 95% of a bisphosphonate active ingredient, and from about 1% to about 99.9%, preferably from 5.00% to about 99.90% of pharmaceutically-acceptable excipients. For risedronate sodium monohydrate or hemipentahydrate, an oral composition comprises, preferably 0.25% to 40%, preferably from about 0.5% to about 30% of a risedronate active ingredient and from about 60% to about 97%, preferably from about 70% to about 99.5% of pharmaceutically-acceptable excipients.

The phrase "safe and effective amount", as used herein means an amount of a compound or composition high enough to significantly positively modify the symptoms and/or condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. The safe and effective amount of active ingredient for use in the method of the invention herein will vary with the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the particular active ingredient being employed, the particular pharmaceutically-acceptable excipients utilized, and like factors within the knowledge and expertise of the attending physician.

The term "risedronate active ingredient" includes the risedronate, risedronate salts, and risedronate esters or any mixture thereof. Any pharmaceutically-acceptable, non-toxic salt or ester of risedronate may be used as the risedronate active ingredient in the dosage forms of the present invention. The salts of risedronate may be acid addition salts, in particular for risedronate the hydrochloride, but any pharmaceutically-acceptable, non-toxic organic or inorganic acid salt may be used. In addition, salts formed with the phosphonic acid group may be used, including,

but not limited to alkali metal salts (K, Na) and alkaline earth metal salts (Ca, Mg) the Ca and Na salts being preferred.

Particularly, other esters of bisphosphonate which are suitable for use as the active ingredient herein are straight chain or branched chain C₁-C₁₈ alkyl esters, including, but not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, amyl, hexyl, heptyl, octyl, nonyl, decyl, lauryl, myristyl, cetyl, and stearyl; straight chain or branched C₂-C₁₈ alkenyl, esters, including but not limited to vinyl, alkyl, undecenyl, and linolenyl; C₃-C₈ cycloalkyl esters, including, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl; aryl ester, including, but not limited to phenyl, toluyl, xylyl, and naphthyl; alicyclic esters, including, but not limited to, menthyl; and aralkyl esters, including, but not limited to benzyl, and phenethyl.

The term "pharmaceutically-acceptable excipients" as used herein includes any physiologically inert, pharmacologically inactive material known to one skilled in the art, which is compatible with the physical and chemical characteristics of the particular active ingredient selected for use. Pharmaceutically-acceptable excipients include, but are not limited to, polymers, resins, plasticizers, fillers, lubricants, binders, disintegrants, solvents, co-solvents, buffer systems, surfactants, preservatives, sweetening agents, flavoring agents, pharmaceutical grade dyes and pigments. All or part of the pharmaceutically-acceptable excipients contained in the pharmaceutical compositions described herein is used to make the film coating which is to be utilized in the novel oral dosage forms described herein.

The term "oral dosage form" as used herein means any pharmaceutical composition intended to be administered to the stomach of an individual via the mouth of said individual

As stated hereinabove, pharmaceutically-acceptable excipients include, but are not limited to polymers, resins, plasticizers, fillers, lubricants, binders, disintegrants, solvents, co-solvents, surfactants, preservatives, sweetener agents, flavoring agents, buffer systems, pharmaceutical-grade dyes and pigments.

The preferred solvent is water.

Flavoring agents among those useful herein include those described in *Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Company, 1990, pp. 1288-1300, incorporated by reference herein. Dyes, or pigments among those useful herein include those described in *Handbook of Pharmaceutical Excipients*, Second Edition pp. 126-134, 1994 by the American Pharmaceutical Association & the Pharmaceutical Press, incorporated by reference herein.

Preferred co-solvents include, but are not limited to, ethanol, glycerin, propylene glycol, polyethylene glycol.

Preferred buffer systems include, but are not limited to potassium acetate, boric carbonic, phosphoric, succinic, malic, tartaric, citric, acetic, benzoic, lactic, glyceric, gluconic, glutaric and
5 glutamic. Particularly preferred are phosphoric, tartaric, citric, and potassium acetate.

Preferred surfactants include, but are not limited to, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene monoalkyl ethers, sucrose monoesters and lanolin esters and ethers.

Preferred preservatives include, but are not limited to, phenol, alkyl esters of parahydroxybenzoic acid, benzoic acid and the salts thereof, boric acid and the salts thereof,
10 sorbic acid and the salts thereof, chlorbutanol, benzyl alcohol, thimerosal, nitromersol, benzalkonium chloride, cetylpyridinium chloride, methyl paraben, and propyl paraben. Particularly preferred are the salts of benzoic acid, cetylpyridinium chloride, methyl paraben and propyl paraben.

Preferred sweeteners include, but are not limited to, sucrose, glucose, saccharin, and
15 aspartame. Particularly preferred are sucrose and saccharin.

Preferred binders include, but are not limited to methycellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, carbomer, povidone, acacia, guar gum, xanthan gum and tragacanth. Particularly preferred are methycellulose, carbomer, xanthan gum, guar gum, povidone and sodium carboxymethylcellulose.

20 Preferred fillers include, but are not limited to lactose, sucrose, maltodextrin, mannitol, starch, and microcrystalline cellulose.

Preferred plasticizers include, but are not limited to polyethylene glycol, propylene glycol, dibutyl phthalate, and castor oil, acetylated monoglycerides, and triacetin.

25 Preferred lubricants include, but are not limited to, magnesium stearate, stearic acid, and talc.

Preferred disintegrants include, but are not limited to, crospovidone, sodium carboxymethyl starch, sodium starch glycolate, sodium carboxymethyl cellulose, alginic acid, clays, and ion exchange resins.

Preferred polymers, include but are not limited to hydroxypropylmethylcellulose (HPMC)
30 alone and/ or in combination with hydroxypropylcellulose (HPC), carboxymethylcellulose, acrylic resins such as Eudragit® RL30D, manufactured by Rohm Pharma GmbH Weiderstadt, West Germany, methylcellulose, ethylcellulose, and polyvinylpyrrolidone or other commercially available film-coating preparations such as Dri-Klear, manufactured by Crompton & Knowles Corp., Mahwah, NJ or Opadry manufactured by Colorcon, West Point, PA.

Other formulations which may be used to administer the bisphosphonate active ingredient. Such formulations include but are not limited to gel formulations as disclosed in WO97/29754 and EP 0 407 344; effervescent formulations as disclosed in WO97/44017; iontophoretic formulations as disclosed in US Patent 5,730,715; and transdermal formulations as
5 disclosed in EP 0 407 345.

The compositions of the present invention allow for greater flexibility in dosage administration and dosing intervals. For example, the compositions of the present invention may be dosed daily, weekly, biweekly or monthly. The safe and effective amount will vary with the particular condition being treated, the age and physical condition of the patient being treated, the
10 severity of the condition, the duration of treatment, and the nature of concurrent therapy.

What is claimed is:

1. A process for selectively producing 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium hemipentahydrate and monohydrate comprising the steps of:

- a) providing an aqueous solution of 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium;
- b) heating the aqueous solution to a temperature from about 45 °C to about 75 °C;
- c) adding a solvent to the aqueous solution characterized in that the solvent is selected from the group consisting of alcohols, esters, ethers, ketones, amides, and nitriles; and
- d) optionally cooling the aqueous solution.

2. The process of Claim 1 characterized in that the aqueous solution is heated to a temperature from about 55 °C to about 75 °C.

3. The process of Claim 1 characterized in that the aqueous solution is heated to about 70 °C and the aqueous solution is not cooled.

4. The process of Claim 1 characterized in that the aqueous solution is heated to a temperature from about 50 °C to about 70 °C.

5. The process of Claim 4 characterized in that the aqueous solution is cooled at a rate of about 0.1 °C to about 2 °C per minute.

6. The process of Claim 1 characterized in that the aqueous solution is heated to about 60 °C then cooled to about 25 °C in about 2 hours.

7. The process of Claim 11 characterized in that the aqueous solution is heated to about 60 °C and maintained at about 60 °C for about 4 hours then cooled to about 25 °C in about 2 hours.

8. The process according to any of the preceding claims characterized in that the solvent is isopropanol.

9. A pharmaceutical composition comprising 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium characterized in that the 3-pyridyl-1-hydroxyethylidene-1,1-

bisphosphonic acid sodium is from about 50% to about 100% hemipentahydrate and from about 50% to about 0% monohydrate.

10. The pharmaceutical composition of Claim 9 characterized in that the 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium is from about 90% to about 100% hemipentahydrate and from about 10% to about 0% monohydrate.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/56983 A3

(51) International Patent Classification⁷: C07F 9/58,
A61K 31/675, A61P 19/00

(21) International Application Number: PCT/US01/03336

(22) International Filing Date: 1 February 2001 (01.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/179,505 1 February 2000 (01.02.2000) US

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, OH
45202 (US).

(72) Inventors: **CAZER, Fredrick, Dana**; 220 Reese Rd.,
Earlville, NY 13312 (US). **PERRY, Gregory, Eugene**;
8 Gerald St., Norwich, NY 13815 (US). **BILLINGS,**
Dennis, Michael; 142 Warner Road, Norwich, NY 13815
(US). **REDMAN-FUREY, Nancy, Lee**; 786 Beaver
Meadow Rd., Smyrna, NY 13464 (US).

(74) Agents: **REED, T., David et al.**; The Procter & Gam-
ble Company, 5299 Spring Grove Avenue, Cincinnati, OH
45217-1087 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility
model), DK, DK (utility model), DM, DZ, EE, EE (utility
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK
(utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(88) Date of publication of the international search report:
7 March 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: SELECTIVE CRYSTALLIZATION OF 3-PYRIDYL-1-HYDROXYETHYLIDENE-1,1-BISPHOSPHONIC ACID
SODIUM AS THE HEMIPENTAHYDRATE OR MONOHYDRATE

(57) Abstract: The present invention discloses 3-pyridyl-1-hydroxyethylidene-1,1-bisphosphonic acid sodium hemipentahydrate and monohydrate, methods of preparing the hemipentahydrate or monohydrate through control of the nucleation temperature and rate of crystallization and pharmaceutical compositions containing one or both of the hydrate forms

WO 01/56983 A3

INTERNATIONAL SEARCH REPORT

Inter Application No
PCT/US 01/03336

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07F9/58 A61K31/675 A61P19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07F A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 132, no. 1, 3 January 2000 (2000-01-03) Columbus, Ohio, US; abstract no. 6284, PERKINS, A. C. ET AL: "Esophageal transit of risedronate cellulose-coated tablet and gelatin capsule formulations" XP002178317 abstract & INT. J. PHARM. (1999), 186(2), 169-175 ,	1-10
A	EP 0 186 405 A (THE PROCTER & GAMBLE CO.) 2 July 1986 (1986-07-02) the whole document --- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

27 September 2001

Date of making of the international search report

12/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

Interr. Application No
PCT/US 91/03336

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

A	EP 0 891 979 A (GADOR S.A.) 20 January 1999 (1999-01-20) the whole document ---	1-10
E	WO 01 15703 A (MERCK & CO., INC.) 8 March 2001 (2001-03-08) claims 1,5,6,15 ---	9,10
E	WO 01 45636 A (MERCK & CO., INC.) 28 June 2001 (2001-06-28) claims 1,21-32 -----	9,10

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. App. No.

PCT/US 01/03336

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 186405	A	02-07-1986	AT 61370 T	15-03-1991
			AU 587001 B2	03-08-1989
			AU 5153485 A	26-06-1986
			CA 1320727 A1	27-07-1993
			DE 3582033 D1	11-04-1991
			DK 599685 A	22-06-1986
			EP 0186405 A2	02-07-1986
			ES 550302 D0	01-08-1987
			ES 8707543 A1	16-10-1987
			ES 557322 D0	01-06-1988
			ES 8802428 A1	16-09-1988
			IE 58749 B	03-11-1993
			IL 77243 A	14-11-1996
			JP 2798666 B2	17-09-1998
			JP 9202794 A	05-08-1997
			JP 2702419 B2	21-01-1998
			JP 7285976 A	31-10-1995
			JP 2568999 B2	08-01-1997
			JP 61210033 A	18-09-1986
			LU 90669 A9	10-01-2001
			NZ 214651 A	26-04-1990
			PH 27078 A	01-02-1993
			US 5583122 A	10-12-1996
			ZA 8509694 A	27-08-1986
EP 891979	A	20-01-1999	EP 0891979 A1	20-01-1999
WO 0115703	A	08-03-2001	AU 6406900 A	26-03-2001
			WO 0115703 A1	08-03-2001
			US 2001018431 A1	30-08-2001
WO 0145636	A	28-06-2001	WO 0145636 A1	28-06-2001